

A CLASS OF LOGARITHMICALLY COMPLETELY MONOTONIC FUNCTIONS RELATING THE q -GAMMA FUNCTION AND APPLICATIONS

KHALED MEHREZ

ABSTRACT. In this paper, the logarithmically complete monotonicity property for a functions involving q -gamma function is investigated for $q \in (0, 1)$. As applications of this results, some new inequalities for the q -gamma function are established. Furthermore, let the sequence r_n be defined by

$$n! = \sqrt{2\pi n} (n/e)^n e^{r_n}.$$

We establish new estimates for Stirling's formula remainder r_n .

Keywords: Completely monotonic functions, Logarithmically completely monotonic functions, q -gamma function, Stirling's formula, Inequalities.

Mathematics Subject Classification (2010): 33D05, 26D07, 26A48

1. Introduction

A real valued function f , defined on an interval I , is called completely monotonic, if f has derivatives of all orders and satisfies

$$(1) \quad (-1)^n f^{(n)}(x) \geq 0, \quad n \in \mathbb{N}_0, \quad x \in I,$$

where $\mathbb{N}$ the set of all positive integers.

A positive function f is said to be logarithmically completely monotonic on an interval I if its logarithm $\log f$ satisfies

$$(-1)^n \left(\log f(x) \right)^{(n)}(x) \geq 0,$$

for all $x \in I$ and $n \in \mathbb{N}$.

Completely monotonic functions have remarkable applications in different branches of mathematics. For instance, they play a role in potential theory, probability theory, physics, numerical and asymptotic analysis, and combinatorics (see [3] and the references given therein).

The q -analogue of the gamma function is defined as

$$(2) \quad \Gamma_q(x) = (1-q)^{1-x} \prod_{j=0}^{\infty} \frac{1-q^{j+1}}{1-q^{j+x}}, \quad 0 < q < 1,$$

and

$$(3) \quad \Gamma_q(x) = (q-1)^{1-x} q^{\frac{x(x-1)}{2}} \prod_{j=0}^{\infty} \frac{1-q^{-(j+1)}}{1-q^{-(j+x)}}, \quad q > 1.$$

The q -gamma function $\Gamma_q(z)$ has the following basic properties:

$$(4) \quad \lim_{q \rightarrow 1^-} \Gamma_q(z) = \lim_{q \rightarrow 1^+} \Gamma_q(z) = \Gamma(z),$$

and

$$(5) \quad \Gamma_q(z) = q^{\frac{(x-1)(x-2)}{2}} \Gamma_{\frac{1}{q}}(z).$$

The q -digamma function ψ_q , the q -analogue of the psi or digamma function ψ is defined for $0 < q < 1$ by

$$(6) \quad \begin{aligned} \psi_q(x) &= \frac{\Gamma'_q(x)}{\Gamma_q(x)} \\ &= -\log(1-q) + \log q \sum_{k=0}^{\infty} \frac{q^{k+x}}{1-q^{k+x}} \\ &= -\log(1-q) + \log q \sum_{k=1}^{\infty} \frac{q^{kx}}{1-q^k}. \end{aligned}$$

For $q > 1$ and $x > 0$, the q -digamma function ψ_q is defined by

$$\begin{aligned} \psi_q(x) &= -\log(q-1) + \log q \left[x - \frac{1}{2} - \sum_{k=0}^{\infty} \frac{q^{-(k+x)}}{1-q^{-(k+x)}} \right] \\ &= -\log(q-1) + \log q \left[x - \frac{1}{2} - \sum_{k=1}^{\infty} \frac{q^{-kx}}{1-q^{-kx}} \right] \end{aligned}$$

From the previous definitions, for a positive x and $q > 1$, we get

$$\psi_q(x) = \frac{2x-3}{2} \log q + \psi_{1/q}(x).$$

Using the Euler-Maclaurin formula, Moak [7] obtained the following q -analogue of Stirling formula

$$(7) \quad \log \Gamma_q(x) \sim \left(x - \frac{1}{2}\right) \log \left(\frac{1-q^x}{1-q}\right) + \frac{\text{Li}_2(1-q^x)}{\log q} + \frac{1}{2} H(q-1) \log q + C_{\hat{q}} + \sum_{k=1}^{\infty} \frac{B_{2k}}{(2k)!} \left(\frac{\log \hat{q}}{\hat{q}^x - 1}\right)^{2k-1} \hat{q}^x P_{2k-3}(\hat{q}^x)$$

as $x \rightarrow \infty$ where $H(\cdot)$ denotes the Heaviside step function, B_k , $k = 1, 2, \dots$ are the Bernoulli numbers,

$$\hat{q} = \begin{cases} q & \text{if } 0 < q < 1 \\ 1/q & \text{if } q > 1 \end{cases}$$

$\text{Li}_2(z)$ is the dilogarithm function defined for complex argument z as [1]

$$(8) \quad \text{Li}_2(z) = - \int_0^z \frac{\log(1-t)}{t} dt, \quad z \notin (0, \infty)$$

P_k is a polynomial of degree k satisfying

$$(9) \quad P_k(z) = (z - z^2) P'_{k-1}(z) + (kz + 1) P_{k-1}(z), \quad P_0 = P_{-1} = 1, \quad k = 1, 2, \dots$$

and

$$C_k = \frac{1}{2} \log(2\pi) + \frac{1}{2} \log \left(\frac{q-1}{\log q} \right) - \frac{1}{24} \log q + \log \left(\sum_{m=-\infty}^{\infty} r^{m(6m+1)} - r^{(2m+1)(3m+1)} \right),$$

where $r = \exp(4\pi^2/\log q)$. It is easy to see that

$$\lim_{q \rightarrow 1} C_q = \frac{1}{2} \log(2\pi), \quad \text{and} \quad \lim_{q \rightarrow 1} \frac{\text{Li}_2(1-q^x)}{\log q} = -x.$$

Stirling's formula

$$(10) \quad n! \sim \sqrt{2\pi n} \left(\frac{n}{e} \right)^n, \quad n \in \mathbb{N}$$

has many applications in statistical physics, probability theory and number theory. Actually, it was first discovered in 1733 by the French mathematician Abraham de Moivre (1667-1754) in the form

$$n! \sim \text{constant} \sqrt{n} \left(\frac{n}{e} \right)^n$$

when he was studying the Gaussian distribution and the central limit theorem. Afterwards, the Scottish mathematician James Stirling (1692-1770) found the missing constant $\sqrt{2\pi}$ when he was trying to give the normal approximation of the binomial distribution.

In 1940 Hummel [4] defined the sequence r_n by

$$(11) \quad n! = \sqrt{2\pi n} (n/e)^n e^{r_n},$$

and established

$$(12) \quad \frac{11}{2} < r_n + \log \sqrt{2\pi} < 1$$

After the inequality (12) was published, many improvements have been given. For example, Robbins [9] established

$$(13) \quad \frac{1}{12n+1} < r_n < \frac{1}{12n}$$

The main aim of this paper is to investigate the logarithmic complete monotonicity property of the function

$$(14) \quad f_{\alpha, \beta}(x; q) = \frac{\Gamma_q(x + \beta) \exp \left(\frac{-\text{Li}_2(1-q^x)}{\log q} \right)}{\left(\frac{1-q^x}{1-q} \right)^{x+\beta-\alpha}}, \quad x > 0,$$

for all reals α, β and q such that $q \in (0, 1)$. As applications of these results, sharp bounds for the q -gamma function are derived. In addition, we present new estimate for for Stirling's formula remainder r_n , (see Corollary 4). Some results are shown to be a generalization of results which were obtained by Chen and Qi [2].

2. logarithmically completely monotonic function related the q -gamma function

In order to study the function defined by (14) we need the following lemma which is considered the main tool to arrive at our results.

Lemma 1. [10] *For every $x, q \in \mathbb{R}_+$, there exists at least one real number $a \in [0, 1]$ such that*

$$(15) \quad \psi_q(x) = \log \left(\frac{1 - q^{x+a}}{1 - q} \right) + \frac{q^x \log q}{1 - q^x} - \left(\frac{1}{2} - a \right) H(q - 1) \log q$$

where $H(\cdot)$ is the Heaviside step function.

Theorem 1. *Let α be a real number. The function $f_{\alpha,1}(q; x)$ is logarithmically completely monotonic on $(0, \infty)$, if and only if $2\alpha \leq 1$.*

Proof. Taking logarithm of $f_{\alpha,1}(x; q)$ leads to

$$(16) \quad \log f_{\alpha,1}(x; q) = \log \Gamma_q(x + 1) - (x + 1 - \alpha) \log \left(\frac{1 - q^x}{1 - q} \right) - \frac{\text{Li}_2(1 - q^x)}{\log q}.$$

Differentiation yields

$$(17) \quad (\log f_{\alpha,1}(x; q))' = \psi_q(x + 1) - \log \left(\frac{1 - q^x}{1 - q} \right) + (1 - \alpha) \frac{\log q \, q^x}{1 - q^x}.$$

From the series expansion

$$\frac{1}{(1 - x)^2} = \sum_{k=0}^{\infty} (k + 1) x^k,$$

for $x \in (0, 1)$ and (6) we get

$$(18) \quad \begin{aligned} (\log f_{\alpha,1}(x; q))'' &= \psi_q'(x + 1) + (1 - \alpha) \frac{q^x}{(1 - q^x)^2} + \log q \frac{q^x}{1 - q^x} \\ &= (\log q)^2 \sum_{k=1}^{\infty} \frac{k q^{k(x+1)}}{1 - q^k} + (1 - \alpha) (\log q)^2 \sum_{k=1}^{\infty} k q^{kx} + \log q \sum_{k=1}^{\infty} q^{kx} \\ &= \sum_{k=1}^{\infty} \frac{\log q \, q^{kx}}{1 - q^k} \Phi_{\alpha,1}(q^k), \end{aligned}$$

where

$$\Phi_{\alpha,1}(y) = y \log y + (1 - \alpha)(1 - y) \log y + (1 - y), \quad y = q^k, \quad k = 1, 2, \dots$$

In order to determine the sign of the function $\Phi_{\alpha,1}(y)$, we have

$$(19) \quad \begin{aligned} \Phi_{\alpha,1}(y) &= y(-\log(1/y) + (1 - \alpha) \log(1/y)(1 - 1/y) + 1/y - 1) \\ &= y \sum_{k=2}^{\infty} \frac{(\log(1/y))^k}{(k - 1)!} \left[\alpha - 1 + \frac{1}{k} \right]. \end{aligned}$$

Therefore, the function $\Phi_{\alpha,1}(y)$ is less than zero if $2\alpha \leq 1$. Thus implies that the function $(\log f_{\alpha,1}(x; q))''$ is completely monotonic on $(0, \infty)$. This can be rewritten as

$$(-1)^n (\log f_{\alpha,1}(x; q))^{(n)} \geq 0, \quad n \geq 2.$$

In particular, $(\log f_{\alpha,1}(x; q))'' \geq 0$, so $(\log f_{\alpha,1}(x; q))'$ is increasing on $(0, \infty)$, and consequently

$$\begin{aligned} (\log f_{\alpha,1}(x; q))^{(1)} &\leq \lim_{x \rightarrow \infty} (\log f_{\alpha,1}(x; q))^{(1)} \\ &= \lim_{x \rightarrow \infty} \left(\psi_q(x+1) - \log \left(\frac{1-q^x}{1-q} \right) + (1-\alpha) \frac{\log q \, q^x}{1-q^x} \right) \\ &= 0. \end{aligned}$$

So $f_{\alpha,1}$ is logarithmically completely monotonic on $(0, \infty)$ if $2\alpha \leq 1$.

Conversely, If the function $f_{\alpha,1}(x; q)$ is logarithmically completely monotonic on $(0, \infty)$, then for all real $x > 0$,

$$(20) \quad (\log f_{\alpha,1}(x; q))' = \psi_q(x+1) - \log \left(\frac{1-q^x}{1-q} \right) + (1-\alpha) \frac{\log q \, q^x}{1-q^x} \leq 0.$$

From the equation (20) and along with the identity

$$(21) \quad \psi_q(x+1) = \psi_q(x) - \frac{q^x \log q}{1-q^x},$$

we have

$$(\log f_{\alpha,1}(x; q))' = \psi_q(x) - \log \left(\frac{1-q^x}{1-q} \right) - \alpha \log q \frac{q^x}{1-q^x} \leq 0,$$

which is equivalent to

$$(22) \quad \psi_q(x) - \log \left(\frac{1-q^x}{1-q} \right) \leq \alpha \frac{\log q \, q^x}{1-q^x}.$$

It is worth mentioning that, Moak [7] proved the following approximation for the q -digamma function

$$\psi_q(x) = \log \left(\frac{1-q^x}{1-q} \right) + \frac{1}{2} \frac{\log q \, q^x}{1-q^x} + O \left(\frac{\log^2 q \, q^x}{(1-q^x)^2} \right)$$

holds for all $q > 0$ and $x > 0$ and so $\psi_q(x) \sim I(x; q)$ on $(0, \infty)$ where

$$(23) \quad I(x; q) = \log \left(\frac{1-q^x}{1-q} \right) + \frac{1}{2} \frac{\log q \, q^x}{1-q^x}$$

Combining (22) and (23) we have

$$\alpha \leq \frac{1}{2}.$$

The proof is complete. ■

Theorem 2. *Let α be a real number. The function $[f_{\alpha,1}(q; x)]^{-1}$ is logarithmically completely monotonic on $(0, \infty)$, if and only if $\alpha \geq 1$.*

Proof. From (19), we conclude that the function $\Phi_{\alpha,1}(y) \geq 0$ if $\alpha \geq 1$ we conclude that

$$(-1)^n \left(\log \frac{1}{f_{\alpha,1}(q; x)} \right)^{(n)} \geq 0$$

for all $x > 0$, $\alpha \geq 1$, $q \in (0, 1)$. and $n \geq 2$. So,

$$\left(\log \frac{1}{f_{\alpha,1}(q; x)} \right)^{(1)} = \log \left(\frac{1 - q^x}{1 - q} \right) - \psi_q(x) + \alpha \log q \frac{q^x}{1 - q^x},$$

is increasing, thus

$$\begin{aligned} \left(\log \frac{1}{f_{\alpha,1}(q; x)} \right)^{(1)} &< \lim_{x \rightarrow \infty} \left(\log \frac{1}{f_{\alpha,1}(q; x)} \right)^{(1)} \\ &= \lim_{x \rightarrow \infty} \left(\log \left(\frac{1 - q^x}{1 - q} \right) - \psi_q(x) + \alpha \log q \frac{q^x}{1 - q^x} \right) \\ &= 0. \end{aligned}$$

Hence, For $\alpha \geq 1$ and $n \in \mathbb{N}$,

$$(-1)^n \left(\log \frac{1}{f_{\alpha,1}(q; x)} \right)^{(n)} \geq 0,$$

on $(0, \infty)$. Now, assume that $\frac{1}{f_{\alpha,1}(q; x)}$ is logarithmically completely monotonic on $(0, \infty)$, by definition, this give us that for all $q \in (0, 1)$ and $x > 0$,

$$\left(\log \frac{1}{f_{\alpha,1}(q; x)} \right)^{(1)} = \log \left(\frac{1 - q^x}{1 - q} \right) - \psi_q(x) + \alpha \log q \frac{q^x}{1 - q^x} \leq 0,$$

which implies that

$$(24) \quad \alpha \geq \frac{1 - q^x}{\log q \, q^x} \left(\psi_q(x) - \log \left(\frac{1 - q^x}{1 - q} \right) \right).$$

In view of Lemma 1 and inequality (24), we see that for all $x > 0$ and $q \in (0, 1)$ there exists at least one real number $a \in [0, 1]$ such that

$$\alpha \geq \frac{1 - q^x}{\log q \, q^x} \left(\log \left(\frac{1 - q^{x+a}}{1 - q^x} \right) + \frac{\log q \, q^x}{1 - q^x} \right),$$

and consequently

$$\alpha \geq 1$$

as $x \rightarrow \infty$. This ends the proof. ■

Theorem 3. *Let α be a real number and $\beta \geq 0$. Then, the function $f_{\alpha,\beta}(x; q)$ is logarithmically completely monotonic function on $(0, \infty)$ if $2\alpha \leq 1 \leq \beta$.*

Proof. Standard calculations lead us to

$$\begin{aligned}
 (\log f_{\alpha,\beta}(x; q))'' &= \psi'_q(x + \beta) + (\beta - \alpha) \frac{q^x}{(1 - q^x)^2} + \log q \frac{q^x}{1 - q^x} \\
 &= (\log q)^2 \sum_{k=1}^{\infty} \frac{k q^{k(x+\beta)}}{1 - q^k} + (\beta - \alpha) (\log q)^2 \sum_{k=1}^{\infty} k q^{kx} + \log q \sum_{k=1}^{\infty} q^{kx} \\
 &= \sum_{k=1}^{\infty} \frac{\log q \cdot q^{kx}}{1 - q^k} \Phi_{\alpha,\beta}(q^k),
 \end{aligned}$$

where

$$\Phi_{\alpha,\beta}(y) = y^\beta \log y + (\beta - \alpha)(1 - y) \log y + (1 - y), \quad y = q^k, \quad k = 1, 2, \dots$$

Thus

$$\Phi_{\alpha,\beta}(y) = y^\beta \left(\sum_{k=2}^{\infty} \frac{(\log(1/y))^k}{(k-1)!} \left[\frac{\beta^k - (\beta-1)^k}{k} + (\beta - \alpha)[(\beta-1)^{k-1} - \beta^{k-1}] \right] \right).$$

From the inequality [2]

$$\beta^k - (\beta-1)^k < k(\beta - \alpha)(\beta^{k-1} - (\beta-1)^{k-1})$$

we conclude that $\Phi_{\alpha,\beta}(y) \leq 0$. This implies that for $n \geq 2$

$$(25) \quad (-1)^n (\log f_{\alpha,\beta}(x; q))^{(n)} \geq 0$$

on $(0, \infty)$ for $2\alpha \leq 1 \leq \beta$. As $(\log f_{\alpha,\beta}(x; q))^{(2)} \geq 0$, it follows that $(\log f_{\alpha,\beta}(x; q))^{(1)}$ is increasing on $(0, \infty)$, and consequently

$$\begin{aligned}
 (\log f_{\alpha,\beta}(x; q))^{(1)} &\leq \lim_{x \rightarrow \infty} (\log f_{\alpha,\beta}(x; q))^{(1)} \\
 &= \lim_{x \rightarrow \infty} \left(\psi_q(x + \beta) - \log \left(\frac{1 - q^x}{1 - q} \right) + (\beta - \alpha) \frac{\log q \cdot q^x}{1 - q^x} \right) \\
 &= 0.
 \end{aligned}$$

In conclusion, (25) is true also $n = 1$, and we conclude that the function $f_{\alpha,\beta}(x; q)$ is logarithmically completely monotonic on $(0, \infty)$ for $2\alpha \leq 1 \leq \beta$. The proof is now completed. $\blacksquare$

3. Inequalities

As applications of the logarithmic complete monotonicity properties of the function (14) which are proved in Theorem 1, Theorem 2 and Theorem 3, we can provide the following inequalities for the q -gamma functions.

Corollary 1. *Let $q \in (0, 1)$, $n \in \mathbb{N}$ and $x_k > 0$ ($1 \leq k \leq n$). Suppose that*

$$\sum_{k=1}^n p_k = 1 \quad (p_k \geq 0).$$

If $2\alpha \leq 1 \leq \beta$, then

(26)

$$\frac{\Gamma_q\left(\sum_{k=1}^n p_k x_k + \beta\right)}{\prod_{k=1}^n \left[\Gamma_q(x_k + \beta)\right]^{p_k}} \leq \frac{\left(\frac{1-q^{\sum_{k=1}^n p_k x_k}}{1-q}\right)^{\sum_{k=1}^n p_k x_k + \beta - \alpha}}{\prod_{k=1}^n \left(\frac{1-q^{x_k}}{1-q}\right)^{p_k(x_k + \beta - \alpha)}} \exp\left(\frac{Li_2\left(1 - q^{\sum_{k=1}^n p_k x_k}\right) - \sum_{k=1}^n p_k Li_2(1 - q^{x_k})}{\log q}\right)$$

Proof. From Theorem 3, $f_{\alpha,\beta}(x; q)$ is logarithmically completely monotonic on the interval $(0, \infty)$, which also implies that the function $f_{\alpha,\beta}(x; q)$ is logarithmically convex. Combining this fact with Jensen's inequality for convex functions yields

$$(27) \quad \log f_{\alpha,\beta}\left(\sum_{k=1}^n p_k x_k; q\right) \leq \sum_{k=1}^n p_k \log f_{\alpha,\beta}(x_k; q).$$

Rearranging (27) can lead to the inequality (26). ■

Corollary 2. Let $q \in (0, 1)$, $n \in \mathbb{N}$ and $x_k > 0$ ($1 \leq k \leq n$). Suppose that

$$\sum_{k=1}^n p_k = 1 \quad (p_k \geq 0).$$

Then, the following inequalities holds

$$(28) \quad \frac{\left(\frac{1-q^{\sum_{k=1}^n p_k x_k}}{1-q}\right)^{\sum_{k=1}^n p_k x_k}}{\prod_{k=1}^n \left(\frac{1-q^{x_k}}{1-q}\right)^{p_k x_k}} \exp\left(\frac{Li_2\left(1 - q^{\sum_{k=1}^n p_k x_k}\right) - \sum_{k=1}^n p_k Li_2(1 - q^{x_k})}{\log q}\right) \\ \leq \frac{\Gamma_q\left(\sum_{k=1}^n p_k x_k + 1\right)}{\prod_{k=1}^n \left[\Gamma_q(x_k + 1)\right]^{p_k}} \\ \leq \frac{\left(\frac{1-q^{\sum_{k=1}^n p_k x_k}}{1-q}\right)^{\sum_{k=1}^n p_k x_k + 1/2}}{\prod_{k=1}^n \left(\frac{1-q^{x_k}}{1-q}\right)^{p_k(x_k + 1/2)}} \exp\left(\frac{Li_2\left(1 - q^{\sum_{k=1}^n p_k x_k}\right) - \sum_{k=1}^n p_k Li_2(1 - q^{x_k})}{\log q}\right)$$

Proof. The right side inequality of (28) follows by inequality (26). From Theorem 2, the function $f_{1,1}(x; q)$ is logarithmically concave. Combining this fact with Jensen's inequality for convex functions we obtain the left side inequality of (28). ■

Corollary 3. Let $q \in (0, 1)$ and a, b be a reals numbers such that $0 < a < b$. Then the following inequalities

(29)

$$\frac{\left(\frac{1-q^b}{1-q}\right)^{b-1}}{\left(\frac{1-q^a}{1-q}\right)^{a-1}} \exp\left(\frac{Li_2(1 - q^b) - Li_2(1 - q^a)}{\log q}\right) \leq \frac{\Gamma_q(b)}{\Gamma_q(a)} \leq \frac{\left(\frac{1-q^b}{1-q}\right)^{b-\frac{1}{2}}}{\left(\frac{1-q^a}{1-q}\right)^{a-\frac{1}{2}}} \exp\left(\frac{Li_2(1 - q^b) - Li_2(1 - q^a)}{\log q}\right)$$

holds.

Proof. From the monotonicity of the functions $f_{\frac{1}{2},1}(q; x)$ and $[f_{1,1}(q; x)]^{-1}$ and the recurrence formula

$$(30) \quad \Gamma_q(x+1) = \frac{1-q^x}{1-q} \Gamma_q(x),$$

we obtain the inequalities (29). ■

Corollary 4. *Let $q \in (0, 1)$ the following inequalities*

$$(31) \quad \left(\frac{1-q^x}{1-q} \right)^x \exp \left(-\frac{Li_2(1-q)}{\log q} \right) \exp \left(\frac{Li_2(1-q^x)}{\log q} \right) \leq \Gamma_q(x+1) \\ \leq \left(\frac{1-q^x}{1-q} \right)^{x+1/2} \exp \left(-\frac{Li_2(1-q)}{\log q} \right) \exp \left(\frac{Li_2(1-q^x)}{\log q} \right)$$

holds for all $x \in [1, \infty)$.

Proof. As the function $f_{1/2,1}(q; x)$ is logarithmically completely monotonic, $f_{1/2,1}(q; x)$ is also decreasing. The following inequality hold true for every $x \geq 1$:

$$(32) \quad f_{1/2,1}(q; x) \leq f_{1/2,1}(q; 1).$$

In addition, as $1/f_{1,1}(q; x)$ is logarithmically completely monotonic, we deduce that $f_{1,1}(q; x)$ is increasing. The following inequality hold true for all $x \geq 1$:

$$(33) \quad f_{1,1}(q; 1) \leq f_{1,1}(q; x).$$

Combining inequalities(32) and (33) we obtain the inequalities (31). ■

In the next Corollary we present new estimates for Stirling's formula remainder r_n .

Corollary 5. *The following inequalities hold true for every integer $n \geq 1$:*

$$(34) \quad e \cdot \left(\frac{n}{e} \right)^n \leq n! \leq e \cdot \sqrt{n} \left(\frac{n}{e} \right)^n,$$

and

$$(35) \quad 1 - \frac{\log 2\pi n}{2} \leq r_n \leq 1 - \frac{\log 2\pi}{2}$$

In each of the above inequalities equality hold if and only if $n = 1$.

Proof. Replacing x by $n \geq 1$ and letting $q \rightarrow 1$ in (31) we get (34), and after some standard computations we obtain inequalities (35). ■

4. Concluding Remarks

1. It is worth mentioning that the inequality (29) when letting q tends to 1, returns to the inequalities [2]

$$(36) \quad \frac{b^{b-1}}{a^{a-1}} e^{a-b} < \frac{\Gamma(a)}{\Gamma(b)} < \frac{b^{b-1/2}}{a^{a-1/2}} e^{a-b},$$

for $b > a > 0$.

2. Let $n = 2$ and $p_k = 1/2$, $k = 1, 2$ in inequalities (28) we obtain the lower bounds for the q -analogue for Gurland's ratio [6] as follows

$$(37) \quad \frac{\left(\frac{1-q^{\frac{x+y}{2}}}{1-q}\right)^{x+y}}{\left(\frac{1-q^x}{1-q}\right)^x \left(\frac{1-q^y}{1-q}\right)^y} \leq \frac{\Gamma_q^2\left(\frac{x+y+2}{2}\right)}{\Gamma_q(x+1)\Gamma_q(y+1)} \exp\left(\frac{\text{Li}_2(1-q^x) + \text{Li}_2(1-q^y) - 2 \text{Li}_2(1-q^{x+y/2})}{\log q}\right) \\ \leq \frac{\left(\frac{1-q^{\frac{x+y}{2}}}{1-q}\right)^{x+y+1}}{\left(\frac{1-q^x}{1-q}\right)^{x+1/2} \left(\frac{1-q^y}{1-q}\right)^{y+1/2}}$$

where $x, y \in (0, \infty)$. In particular, let q tends to 1 in (37) we get

$$(38) \quad \frac{\left(\frac{x+y}{2}\right)^{x+y}}{x^x y^y} \leq \frac{\Gamma^2\left(\frac{x+y+2}{2}\right)}{\Gamma(x+1)\Gamma(y+1)} \leq \frac{\left(\frac{x+y}{2}\right)^{x+y+1}}{x^{x+1/2} y^{y+1/2}}.$$

The left hand side inequalities of (38) has proved first by Mortici [8] and the right hand side of inequalities (38) is new.

REFERENCES

- [1] M. Abramowitz, C.A. Stegun, Handbook of Mathematical Functions with Formulas, Graphs, Mathematical Tables 7th Printing, in: Applied Mathematics Series, vol. 55, National Bureau of Standards, Washington, DC, 1964.
- [2] C.-P. Chen, F. Qi, Logarithmically completely monotonic functions relating to the gamma function, J. Math. Anal. Appl. 321 (2006) 405411.
- [3] C.-P. Chen and F. Qi, Logarithmically completely monotonic ratios of mean values and an application, Glob. J. Math. Math. Sci. 1 (2005) 7176.
- [4] P.M. Hummel, A note on Stirling's formula, Am. Math. Mon. 47 (1940) 9799.
- [5] C. Krattenthaler, H.M. Srivastava, Summations for basic hypergeometric series involving a q -analogue of the digamma function, Computers and Mathematics with Applications 32 (2) (1996) 7391.
- [6] M. Merkle, Gurland's ratio for the gamma function, Comput. Math. Appl. 49 (2005) 389406.
- [7] D. S. Moak, The q -analogue of Stirlings formula, Rocky Mountain J. Math. 14 (1984) 403412.
- [8] C. Mortici, Completely monotonic functions associated with gamma function and applications, Carpathian J. Math. (25), 2009, p. 186-191.
- [9] H. Robbins, A remark on Stirling's formula, Am. Math. Mon. 62 (1955) 2629.
- [10] A. Salem, Ceratin class of approximation for the q -digamma function, Rocky Mountain Journal of Mathematics, (To appear).

KHALED MEHREZ. DÉPARTEMENT DE MATHÉMATIQUES ISSAT KASSERINE, TUNISIA.

E-mail address: k.mehrez@yahoo.fr